

THE ROKHLIN PROPERTY FOR AUTOMORPHISMS ON SIMPLE C^* -ALGEBRAS

JIAJIE HUA

ABSTRACT. Let $\mathcal{A}$ be the class of unital separable simple amenable C^* -algebras A which satisfy the Universal Coefficient Theorem for which $A \otimes M_{\mathfrak{p}}$ has tracial rank zero for some supernatural number $\mathfrak{p}$ of infinite type. Let $A \in \mathcal{A}$ and let α be an automorphism of A . Suppose that α has the tracial Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$, we show that $A \rtimes_{\alpha} \mathbb{Z} \in \mathcal{A}$.

1. INTRODUCTION

Alan Connes introduced the Rokhlin property in ergodic theory to operator algebras ([2]). Several versions of the Rokhlin property for automorphism on C^* -algebra have been studied (for example [7],[15],[14],[4]). Given a unital C^* -algebra A and an automorphism α of A , one may view the pair (A, α) as a non-commutative dynamical system. To study its dynamical structure, it is natural to introduce the notion of Rokhlin property. Let A be a unital simple AT-algebra (direct limits of circle algebras) with real rank zero and let α be an automorphism of A . In several cases, Kishimoto showed that if α is approximately inner and has a Rokhlin property, then the crossed product $A \rtimes_{\alpha} \mathbb{Z}$ is again a unital simple AT with real rank zero ([8],[7]).

In [10], H. Lin proved the following result: Let A be a unital separable simple C^* -algebra with tracial rank zero and let α be an automorphism of A . Suppose that α has a tracial Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$. Then $A \rtimes_{\alpha} \mathbb{Z}$ has tracial rank zero. Thus H. Lin generalizes the Kishimoto's results on unital simple AT-algebras with real rank zero to C^* -algebras with tracial rank zero.

In this paper, we generalize Lin's result as the following: Let $\mathcal{A}$ be the class of unital separable simple amenable C^* -algebras A which satisfy the Universal Coefficient Theorem for which $A \otimes M_{\mathfrak{p}}$ has tracial rank zero for some supernatural number $\mathfrak{p}$ of infinite type. Let $A \in \mathcal{A}$ and let α be an

2000 Mathematics Subject Classification. Primary 46L55; Secondary 46L35, 46L40.

Key words and phrases. crossed products; tracial rank zero; tracial Rokhlin property.

The author was supported the National Natural Science Foundation of China (Nos. 10771069, 10671068, 10771161).

automorphism of A . Suppose that α has the tracial Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$. Then $A \rtimes_\alpha \mathbb{Z} \in \mathcal{A}$.

This paper is organized as follows. In Section 2 we introduce notation on C^* -algebras. In Section 3, we prove that, under our hypotheses, $A \rtimes_\alpha \mathbb{Z} \in \mathcal{A}$.

2. NOTATION

We will use the following convention:

(1) Let A be a C^* -algebra, let $a \in A$ be a positive element and let $p \in A$ be a projection. We write $[p] \leq [a]$ if there is a projection $q \in \overline{aAa}$ and a partial isometry $v \in A$ such that $v^*v = p$ and $vv^* = q$.

(2) Let A be a C^* -algebra. We denote by $\text{Aut}(A)$ the automorphism group of A . If A is unital and $u \in A$ is a unitary, we denote by adu the inner automorphism defined by $\text{adu}(a) = u^*au$ for all $a \in A$.

(3) Let $x \in A$, $\varepsilon > 0$ and $\mathcal{F} \subset A$. We write $x \in_\varepsilon \mathcal{F}$, if $\text{dist}(x, \mathcal{F}) < \varepsilon$, or there is $y \in \mathcal{F}$ such that $\|x - y\| < \varepsilon$.

(4) Let A be a unital C^* -algebra and $T(A)$ the compact convex set of tracial states of A .

(5) A unital C^* -algebra is said to have real rank zero, written $\text{RR}(A) = 0$, if the set of invertible self-adjoint elements is dense in self-adjoint elements of A .

(6) we say the order on projection over a unital C^* -algebra A is determined by traces, if for any two projections $p, q \in A$, $\tau(p) < \tau(q)$ for all $\tau \in T(A)$ implies that p is equivalent to a projection $p' \leq q$.

Definition 2.1. [9] *Let A be a unital simple C^* -algebra. Then A is said to have tracial (topological) rank zero if for every $\varepsilon > 0$, every finite set $\mathcal{F} \subset A$, and every nonzero positive element $a \in A$, there exists a finite dimensional C^* -subalgebra $B \subset A$ with $\text{id}_B = p$ such that:*

- (1) $\|px - xp\| < \varepsilon$ for all $x \in \mathcal{F}$.
- (2) $pap \in_\varepsilon B$ for all $x \in \mathcal{F}$.
- (3) $[1 - p] \leq [a]$.

If A has tracial rank zero, we write $\text{TR}(A) = 0$.

Definition 2.2. (The Jiang-Su algebra [6]) *Denote by $\mathcal{Z}$ the Jiang-Su algebra of unital infinite dimensional simple C^* -algebra which is an inductive limit of prime dimension drop algebras with a unique tracial state, $(K_0(\mathcal{Z}), K_0(\mathcal{Z})_+, [1_{\mathcal{Z}}]) = (\mathbb{Z}, \mathbb{N}, 1)$ and $K_1(\mathcal{Z}) = 0$.*

Definition 2.3. (A classifiable class of unital separable simple amenable C^* -algebra) *Denote by $\mathcal{N}$ the class of all unital separable amenable C^* -algebras which satisfy the Universal Coefficient Theorem.*

For a supernatural number $\mathbf{p}$, denote by $M_{\mathbf{p}}$ the UHF algebra associated with $\mathbf{p}$ (see [3]).

Let $\mathcal{A}$ denote the class of all unital separable simple amenable C^* -algebras A in $\mathcal{N}$ for which $A \otimes M_p$ has tracial rank zero for some supernatural number p of infinite type.

Definition 2.4. Let A be a unital stably finite separable simple amenable C^* -algebra.

By $\text{Ell}(A)$ we mean the following:

$$(K_0(A), K_0(A)_+, [1_A], K_1(A), T(A), r_A),$$

where $r_A : T(A) \rightarrow S_{[1_A]}(K_0(A))$ is a surjective continuous affine map such that $r_A(\tau)([p]) = \tau(p)$ for all projections $p \in A \otimes M_k$ for $k = 1, 2, \dots$.

Suppose that B is another stably finite unital separable simple C^* -algebra. A map $\Lambda : \text{Ell}(A) \rightarrow \text{Ell}(B)$ is said to be a homomorphism if Λ gives an order homomorphism $\lambda_0 : K_0(A) \rightarrow K_0(B)$ such that $\lambda_0 : K_0(A) \rightarrow K_0(B)$ such that $\lambda_0([1_A]) = [1_B]$, a homomorphism $\lambda_1 : K_1(A) \rightarrow K_1(B)$, a continuous affine map $\lambda'_\rho : T(B) \rightarrow T(A)$ such that

$$\lambda'_\rho(\tau)(p) = r_B(\tau)(\lambda_0([p]))$$

for all projection in $A \otimes M_k$ for $k = 1, 2, \dots$ and all $\tau \in T(B)$.

We say that such Λ is an isomorphism, if λ_0 and λ_1 are isomorphisms and λ'_ρ is an affine homeomorphism. In this case, there is an affine homeomorphism $\lambda_\rho : T(A) \rightarrow T(B)$ such that $\lambda_\rho^{-1} = \lambda'_\rho$.

Theorem 2.5. (Corollary 11.9 of [11]) Let $A, B \in \mathcal{A}$. Then

$$A \otimes \mathcal{Z} \cong B \otimes \mathcal{Z}$$

if

$$\text{Ell}(A \otimes \mathcal{Z}) \cong \text{Ell}(B \otimes \mathcal{Z}).$$

Definition 2.6. Recall that a C^* -algebra A is said to be $\mathcal{Z}$ -stable if $A \otimes \mathcal{Z} \cong A$. Denote by $\mathcal{A}_{\mathcal{Z}}$ the class of $\mathcal{Z}$ -stable C^* -algebras in $\mathcal{A}$.

Corollary 2.7. Let A and B be two unital separable amenable simple C^* -algebras in $\mathcal{A}_{\mathcal{Z}}$. Then $A \cong B$ if and only if

$$\text{Ell}(A) \cong \text{Ell}(B).$$

3. MAIN RESULT

We recall the definition of the tracial Rokhlin property.

Definition 3.1. ([13]) Let A be a unital C^* -algebra and let $\alpha \in \text{Aut}(A)$. We say α has the tracial Rokhlin property if for every $\varepsilon > 0$, every $n \in \mathbb{N}$, every nonzero positive element $a \in A$, every finite set $\mathcal{F} \subset A$, there are the mutually orthogonal projections $e_0, e_1, \dots, e_n \in A$ such that:

- (1) $\|\alpha(e_j) - e_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$.
- (2) $\|e_j b - b e_j\| < \varepsilon$ for $0 \leq j \leq n$ and all $b \in \mathcal{F}$.
- (3) With $e = \sum_{j=0}^n e_j$, $[1 - e] \leq [a]$.

The following result is Lemma 1.4 of [13].

Lemma 3.2. *Let A be a stably finite simple unital C^* -algebra such that $RR(A) = 0$ and the order on projections over A is determined by traces. Let $\alpha \in \text{Aut}(A)$. Then α has the tracial Rokhlin property if and only if for every finite set $\mathcal{F} \subset A$, every $\varepsilon > 0$, and every $n \in \mathbb{N}$, there are mutually orthogonal projections $e_0, e_1, \dots, e_n \in A$ such that:*

- (1) $\|\alpha(e_j) - e_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$.
- (2) $\|e_j b - b e_j\| < \varepsilon$ for $0 \leq j \leq n$ and all $b \in \mathcal{F}$.
- (3) With $e = \sum_{j=0}^n e_j$, we have $\tau(1-e) < \varepsilon$ for all $\tau \in T(A)$.

Theorem 3.3. *Let $A \in \mathcal{A}$ be a unital separable simple amenable C^* -algebra, and let α be an automorphism of A . Suppose that α has the tracial Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$. Then $A \rtimes_{\alpha} \mathbb{Z} \in \mathcal{A}$.*

Proof. Since $A \in \mathcal{A}$, we have $TR(A \otimes M_p) = 0$ for some supernatural number p of infinite type. Hence $A \otimes M_p$ have real rank zero, stable rank one and the order on projections over $A \otimes M_p$ is determined by traces (see [9]).

For every $\varepsilon > 0$, every $n \in \mathbb{N}$, every finite set $F \subset A \otimes M_p$, Without loss of generality, we may assume that there exist a finite set $F_A \subset A$ and $F_{M_p} \subset M_p$ such that $F = F_A \otimes F_{M_p}$. We can also find a positive element a of A such that $\tau(a) < \varepsilon$ for all $\tau \in T(A)$.

Since α has the tracial Rokhlin property, there exist mutually orthogonal projections

$$e_0, e_1, \dots, e_n \in A$$

such that:

- (1) $\|\alpha(e_j) - e_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$.
- (2) $\|e_j b - b e_j\| < \varepsilon$ for $0 \leq j \leq n$ and all $b \in F_A$.
- (3) With $e = \sum_{j=0}^n e_j$, $[1-e] \leq [a]$.

Set $e'_j = e_j \otimes 1_{M_p}$ for $0 \leq j \leq n$. Then e'_j , $0 \leq j \leq n$, are mutually orthogonal projections in $A \otimes M_p$.

Then

- (1) $\|(\alpha \otimes \text{id})(e'_j) - e'_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$.
- (2) $\|e'_j b - b e'_j\| < \varepsilon$ for $0 \leq j \leq n$ and all $b \in F$.
- (3) With $e' = \sum_{j=0}^n e'_j$, $\tau(1-e') \leq \tau(a \otimes 1_{M_p}) < \varepsilon$ for all $\tau \in T(A \otimes M_p)$.

So $\alpha \otimes \text{id} \in \text{Aut}(A \otimes M_p)$ has the tracial Rokhlin property by Lemma 3.2.

From assumption, there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$, then $[(\alpha \otimes \text{id})^J] = [\text{id}_A \otimes \text{id}_{M_p}]$ in $KL(A \otimes M_p, A \otimes M_p)$.

Apply Theorem 4.4 of [10], we get $TR((A \otimes M_p) \rtimes_{\alpha \otimes \text{id}} \mathbb{Z}) = 0$.

Next we prove that $(A \otimes M_p) \rtimes_{\alpha \otimes \text{id}} \mathbb{Z} \cong (A \rtimes_{\alpha} \mathbb{Z}) \otimes M_p$.

Let u be the unitary implementing the action of α in the crossed product C^* -algebra $A \rtimes_{\alpha} \mathbb{Z}$.

Set of elements which have form $\sum_{i=-m}^m (\sum_{j=0}^{s_i} a_{ij} \otimes m_{ij})(u \otimes 1_{M_p})^i$ is dense in $(A \otimes M_p) \rtimes_{\alpha \otimes \text{id}} \mathbb{Z}$. We define a map on dense elements of $(A \otimes M_p) \rtimes_{\alpha \otimes \text{id}} \mathbb{Z}$

by

$$\varphi : \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} \otimes m_{ij} \right) (u \otimes 1_{M_P})^i \rightarrow \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} u^i \right) \otimes m_{ij}.$$

Note

$$\begin{aligned} & \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} \otimes m_{ij} \right) (u \otimes 1_{M_P})^i \\ &= \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} \otimes m_{ij} \right) (u^i \otimes 1_{M_P}^i) \\ &= \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} \otimes m_{ij} \right) (u^i \otimes 1_{M_P}) \\ &= \sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} u^i \right) \otimes m_{ij}. \end{aligned}$$

then φ is a $*$ -homomorphism, we can extend φ to $(A \otimes M_P) \rtimes_{\alpha \otimes id} \mathbb{Z}$.

So φ is a norm invariant $*$ -homomorphism, φ is injective, Set of all elements like the form $\sum_{i=-m}^m \left(\sum_{j=0}^{s_i} a_{ij} u^i \right) \otimes m_{ij}$ is dense in $(A \rtimes_{\alpha} \mathbb{Z}) \otimes M_P$. We can extend φ as an isomorphism $(A \otimes M_P) \rtimes_{\alpha \otimes id} \mathbb{Z} \rightarrow (A \rtimes_{\alpha} \mathbb{Z}) \otimes M_P$. then φ is a $*$ -isomorphism.

So $TR((A \rtimes_{\alpha} \mathbb{Z}) \otimes M_P) = 0$, thus $A \rtimes_{\alpha} \mathbb{Z} \in \mathcal{A}$. $\square$

Next we recall the definition of the Rokhlin property.

Definition 3.4. ([8]) Let A be a unital simple C^* -algebra and $\alpha \in \text{Aut}(A)$. We say α has the Rokhlin property if for every $\varepsilon > 0$, every $n \in \mathbb{N}$, every finite set $F \subset A$, there exist the mutually orthogonal projections

$$e_0, e_1, \dots, e_{n-1}, f_0, f_1, \dots, f_n \in A$$

such that:

- (1) $\|\alpha(e_j) - e_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-2$ and $\|\alpha(f_j) - f_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$.
- (2) $\|e_j a - a e_j\| < \varepsilon$ for $0 \leq j \leq n-1$ and all $a \in F$, $\|f_j a - a f_j\| < \varepsilon$ for $0 \leq j \leq n$ for all $a \in F$.
- (3) $\sum_{j=0}^{n-1} e_j + \sum_{j=0}^n f_j = 1$.

If A is a simple unital C^* -algebra with real rank zero, stable rank one, and has weakly unperforated $K_0(A)$, the Rokhlin property implies the tracial Rokhlin property (see [12]).

Corollary 3.5. Let $A \in \mathcal{A}$ be a unital separable simple amenable C^* -algebra, and let α be an automorphism of A . Suppose that α has the Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [id_A]$ in $KL(A, A)$. Then $A \rtimes_{\alpha} \mathbb{Z} \in \mathcal{A}$.

Proof. Using the similar proof of Theorem 3.3, we can know that $\alpha \otimes \text{id}$ has the Rokhlin property. Since $TR(A \otimes M_P) = 0$, by Theorem 1.12 of [13], $\alpha \otimes \text{id}$ has the tracial Rokhlin property. So we can complete the proof as the same as Theorem 3.3. $\square$

The following definition is a stronger version of the Rokhlin property.

Definition 3.6. *Let A be a unital simple C^* -algebra and $\alpha \in \text{Aut}(A)$. We say α has the cyclic Rokhlin property if for every $\varepsilon > 0$, every $n \in \mathbb{N}$, every finite set $F \subset A$, there exist the mutually orthogonal projections*

$$e_0, e_1, \dots, e_{n-1}, f_0, f_1, \dots, f_n \in A$$

such that:

- (1) $\|\alpha(e_j) - e_{j+1}\| < \varepsilon$ for $0 \leq j \leq n-1$ and $e_n = e_0$. $\|\alpha(f_j) - f_{j+1}\| < \varepsilon$ for $0 \leq j \leq n$ and $f_{n+1} = f_0$.
- (2) $\|e_j a - a e_j\| < \varepsilon$ for $0 \leq j \leq n-1$ and for $a \in F$. $\|f_j a - a f_j\| < \varepsilon$ for $0 \leq j \leq n$ and all $a \in F$.
- (3) $\sum_{j=0}^{n-1} e_j + \sum_{j=0}^n f_j = 1$.

The only difference between the Rokhlin property and the cyclic Rokhlin property is that in condition (1) we require that $\|\alpha(e_{n-1}) - e_0\| < \varepsilon$ and $\|\alpha(f_n) - f_0\| < \varepsilon$.

Corollary 3.7. *Let $A \in \mathcal{A}_{\mathcal{Z}}$ be a unital separable simple amenable C^* -algebra, and let α be an automorphism of A . Suppose that α has the cyclic Rokhlin property. Suppose also that there is an integer $J \geq 1$ such that $[\alpha^J] = [\text{id}_A]$ in $KL(A, A)$. Then $A \rtimes_{\alpha} \mathbb{Z} \in \mathcal{A}_{\mathcal{Z}}$.*

Proof. By Theorem 4.4 of [5], if A is $\mathcal{Z}$ -stable, α has the cyclic Rokhlin property, then $A \rtimes_{\alpha} \mathbb{Z}$ is also $\mathcal{Z}$ -stable, we can get the result from Corollary 3.5. $\square$

Remark 3.8. *If A is a unital separable simple C^* -algebra with locally finite decomposition rank which satisfies the UCT. Suppose the projections of A separate traces, since $A \otimes M_P$ is approximate divisible, apply Theorem 1.4(e) of [1], $A \otimes M_P$ has real rank zero. But $A \otimes M_P$ has locally finite decomposition rank and is $\mathcal{Z}$ -stable, by Theorem 2.1 of [16], we have $TR(A \otimes M_P) = 0$. This C^* -algebras class is larger than class of C^* -algebras which have tracial rank zero.*

REFERENCES

- [1] B. Blackadar, A. Kumjian and M. Rørdam, *Approximately central matrix units and the structure of noncommutative tori*, K-theory **6** (1992), 267-284.
- [2] A. Connes, *Outer conjugacy class of automorphisms of factors*, Ann. Sci. Ecole Norm. Sup. **8**(1975), 383-420.
- [3] J. Dixmier, *On some C^* -algebras considered by Glimm*, J. Funct. Anal. **1**(1967), 182-203.
- [4] R. Herman and A. Ocneanu, *Stability for integer actions on UHF- C^* -algebras*, J. Funct. Anal. **59**(1984), 132-144.

- [5] I. Hirshberg and W. Winter, *Rokhlin actions and self-absorbing C^* -algebras*, Pacific J. Math. 233, no. 1 (2007), 125–143.
- [6] X. Jiang and H. Su, *On a simple unital projectionless C^* -algebra*, American J. Math. 121 (1999), no. 2, 359–413.
- [7] A. Kishimoto, *The Rohlin property for automorphisms of UHF algebras*, J. reine. angew. Math. **465**(1995), 183–196.
- [8] A. Kishimoto, *Automorphisms of AT-algebras with the Rohlin property*, Operator Theory. 40(1998),277–294.
- [9] H. Lin, *Tracial topological ranks of C^* -algebra*, Proc. London Math. Soc. **83**(2001), 199–234.
- [10] H. Lin, *The Rokhlin property for automorphisms on simple C^* -algebras*, arXiv:math/0602513v2[math.OA] 27 Jan 2010.
- [11] H. Lin, *Asymptotically unitary equivalence and classification of simple amenable C^* -algebras*, arXiv: 0806.0636(2008).
- [12] H. Osaka and N. C. Phillips, *Furstenberg transformations on irrational rotation algebras*, in preprint.
- [13] H. Osaka and N. C. Phillips, *Stable and real rank for crossed products by automorphisms with the tracial Rokhlin property*. Ergodic Theory Dynam. Systems 26 (2006), no. 5, 1579–1621.
- [14] N. C. Phillips, *Crossed products by finite cyclic group actions with the tracial Rokhlin property*, preprint.
- [15] M. Rørdam, *Classification of certain infinite simple C^* -algebras*, J. Funct. Anal. **131**(1995),415–458.
- [16] W. Winter, *Simple C^* -algebras with locally finite decomposition rank*, J. Funct. Anal. **243**(2007), 394–425.

DEPARTMENT OF MATHEMATICS, TONGJI UNIVERSITY, SHANGHAI 200092,
P.R.CHINA.

E-mail address: huajiajie2006@hotmail.com